

Optical binding of particles with or without the presence of a flat dielectric surface

P. C. Chaumet and M. Nieto-Vesperinas

*Instituto de Ciencia de Materiales de Madrid, Consejo Superior de Investigaciones Científicas, Campus de Cantoblanco
Madrid 28049, Spain*

Optical fields can induce forces between microscopic objects, thus giving rise to new structures of matter. We study theoretically these optical forces between two spheres, either isolated in water, or in presence of a flat dielectric surface. We observe different behavior in the binding force between particles at large and at small distances (in comparison with the wavelength) from each other. This is due to the great contribution of evanescent waves at short distances. We analyze how the optical binding depends of the size of the particles, the material composing them, the wavelength and, above all, on the polarization of the incident beam. We also show that depending on the polarization, the force between small particles at small distances changes its sign. Finally, the presence of a substrate surface is analyzed showing that it only slightly changes the magnitudes of the forces, but not their qualitative nature, except when one employs total internal reflection, case in which the particles are induced to move together along the surface.

I. INTRODUCTION

Some time ago, it was demonstrated that optical fields can produce forces on neutral particles,^{1,2} since then this mechanical action has been used in optical tweezers³ and more recently in optical force microscopy,^{4,5} as well as in manipulating molecules⁶ and dielectric spheres.⁷⁻⁹ In addition, the possibility of binding objects¹⁰ through optical forces and thus create microstructures, either in or off resonant conditions,¹¹⁻¹³ were pointed out.

In this paper we wish to undertake a detailed study on optical forces on neutral particles, based on a rigorous analysis that we have carried out¹⁴⁻¹⁶ in a full three-dimensional configuration by using the coupled dipole method (CDM) of Purcell and Pennypacker¹⁷ via the Maxwell's stress tensor.¹⁸ Specifically, we study the forces induced by light between two spheres, either isolated in solution, or in the presence of a flat dielectric surface. We shall monitor the nature, either attractive or repulsive, of the light induced force between the spheres, according to the wavelength, polarization of the incident wave, and the size and composition of the spheres.

In Section II we outline the calculation method employed to determine the optical binding forces, then, in Section III we present results for spheres either isolated in water (III A) or suspended in this liquid in the presence of a flat dielectric interface (III B).

II. METHOD USED FOR COMPUTING THE OPTICAL BINDING

In a previous article¹⁴ we showed the possibility to compute the optical forces on a sphere with the coupled dipole method (CDM).¹⁷ For the computation of the optical binding between particles, we now use the same procedure, thus we shall next only outline the main equations and the changes introduced in them to address the presence of multiple objects.

Let K objects be above a flat dielectric surface. Each object is discretized into N_k subunits, with $k = 1, \dots, K$. Following the procedure of Ref. [17], the field at the (i, k) th subunit, namely, the i th subunit of the k th object, can be written as:

$$\mathbf{E}(\mathbf{r}_i^k, \omega) = \mathbf{E}_0(\mathbf{r}_i^k, \omega) + \sum_{l=1}^K \sum_{j=1}^{N_l} [\mathbf{S}(\mathbf{r}_i^k, \mathbf{r}_j^l, \omega) + \mathbf{T}(\mathbf{r}_i^k, \mathbf{r}_j^l, \omega)] \alpha_j^l \mathbf{E}(\mathbf{r}_j^l, \omega). \quad (1)$$

where α_j^l is the polarizability of the (j, l) subunit, $\mathbf{T}$ is the linear response to a dipole in free space,¹⁹ and $\mathbf{S}$ represents the linear response of a dipole in the presence of a surface.^{20,21} The value of the electric field at each subunit position is obtained by solving the linear system Eq. (1) written for all subunits, so that the size of the system to solve is $\prod_{k=1}^K N_k$. Once the electric field is obtained, the component of the total averaged force on the (i, k) th subunit can be deduced from both the field and its derivative at its position $\mathbf{r}_i^k$:¹⁵

$$F_u(\mathbf{r}_i^k) = (1/2) \Re e \sum_{v=1}^3 \left(p_v(\mathbf{r}_i^k, \omega) \frac{\partial E_v^*(\mathbf{r}_i^k, \omega)}{\partial u} \right), \quad (u=1, 2, 3). \quad (2)$$

where u, v , stand for either x, y, z , and $\mathbf{p}(\mathbf{r}_i^k, \omega)$ is the electric dipole of the (i, k) th subunit due to the incident field and all the other subunits. Notice, that the derivative of the field can be obtained from the derivative of Eq. (1).¹⁴ Then the following relation can be written:

$$\mathbf{F}^k = \sum_{i=1}^{N_k} \mathbf{F}(\mathbf{r}_i^k) \quad (3)$$

where $\mathbf{F}^k$ is the total force on the k th object due to both the incident field and the multiple interaction with the

surface and the other $(K - 1)th$ objects. If the object kth is a sphere small compared to the wavelength, the dipole approximation can be done, hence $N_k = 1$. We also remark that, in what follows, when we represent the normalized force, this means $\mathbf{F}/(4\pi\epsilon_0|\mathbf{E}_i|^2)$ where ϵ_0 is the permittivity of vacuum and $|\mathbf{E}_i|^2$ denotes the intensity of the incident beam.

III. RESULTS AND DISCUSSION

In Fig. 1 we represent the more complex geometry that we shall consider in this work. Two spheres (either dielectric or metallic) are embedded in water ($\epsilon_w = 1.69$). Illumination with an incident plane wave takes place in the XZ plane at an angle θ of incidence. When a dielectric flat surface at $z = 0$ is used, we consider it separating glass ($\epsilon_s = 2.25$) at $z < 0$ from water ($z > 0$).




FIG. 1. The most complex geometry considered in this paper: two spheres of radius r on a dielectric flat surface. The spheres are embedded in water with $\epsilon_w = 1.69$, and the relative permittivity of the surface is $\epsilon_s = 2.25$. The incident wave vector $\mathbf{k}_0$ is in the XZ plane, and θ is the angle of incidence.

A. Particles in water

In this section we do not address yet the presence of the surface ($\epsilon_s = \epsilon_w = 1.69$), i.e. $\mathbf{S}(\mathbf{r}_i^k, \mathbf{r}_j^l, \omega) = 0$ in Eq. (1), and the angle of incidence is $\theta = 0^\circ$. Even in the absence of surface, we make reference to the polarization and thus we shall always use the terms p -polarization and s -polarization when the electric field vector is in the XZ plane and along the Y -axis, respectively.




FIG. 2. Normalized force in the X -direction on sphere B versus distance x between the centers of the spheres. Both spheres are of glass ($\epsilon_1 = \epsilon_2 = 2.25$), with $r = 10\text{nm}$. The angle of incidence of the illuminating plane wave is $\theta = 0^\circ$ and the wavelength $\lambda = 632.8\text{nm}$ in vacuum. The full line corresponds to p -polarization, and the dashed line represents s -polarization. a) Force for short distances between the spheres, the symbol $+$ ($\times$) corresponds to the values from the non-retarded approximation for p -polarization (s -polarization). b) Force in far field, the symbol $+$ ($\times$) represents the values from the non-retarded approximation in far field for p -polarization (s -polarization).

We begin with the most simple case, i.e., the radius r of the two particles is small compared to the wavelength employed. As previously said, we then use the dipole approximation. We study, first, the case of two identical spheres with $\epsilon_1 = \epsilon_2 = 2.25$, radius $r = 10\text{nm}$ at a wavelength $\lambda = 632.8\text{nm}$ in vacuum. Fig. 2 represents the force along the X -direction on the sphere B at different positions of this sphere on the X -axis. The sphere A remaining fixed. We have only plotted the force exerted on sphere B, since by symmetry, the force along the X -axis on sphere A is the opposite to that on B. We observe two facts: first, the oscillation of the force when the spheres are far from each other, and, second, the strong force, either attractive or repulsive, when the spheres are very close from each other, depending on the polarization. For a better understanding of the physical process, using Eq. (1) and its derivative for two dipolar objects, we can analytically determine, through Eq. (2), the force on the spheres. Then, on using the fact that there is a plane wave in the Z -direction (the incident wave is $E_{0i}e^{ik_0z}$, where $i = x$ or y depending of the polarization of the incident field) the force on the second sphere can be written as:

$$F_x(\mathbf{r}_2) = 1/2\Re e \left(\alpha_2 E_i(\mathbf{r}_2, \omega) \alpha_1^* E_i^*(\mathbf{r}_1, \omega) \frac{\partial}{\partial x} T_{ii}^*(\mathbf{r}_2, \mathbf{r}_1, \omega) \right) \quad (4)$$

where $i = x$ for p -polarization and $i = y$ for s -polarization of the incident field. Notice that to obtain

the force on sphere A, the indices 1 and 2 must be permuted. But, even in this simple case, the exact analytical solution of Eq. (4) is not easy to interpret. Hence, we make in Eq. (1) the approximation that the term $\mathbf{T}(\mathbf{r}_i^k, \mathbf{r}_j^l, \omega)\alpha_j^l$ is smaller than 1 (we will discuss further this approximation). Now, if we used the hypothesis that the two spheres are identical ($\alpha_1 = \alpha_2$), Eq. (4) becomes :

$$F_x(\mathbf{r}_2) = 1/2|\alpha_1 E_{0i}|^2 \Re \left(\frac{\partial}{\partial x} T_{ii}(\mathbf{r}_2, \mathbf{r}_1, \omega) \right). \quad (5)$$

At short distance we can make the non retarded approximation ($k_0 = 0$) and, as shown in the appendix, we have that $F_x(\mathbf{r}_2) = -3|\alpha_1 E_{0x}|^2/a^4$ in p -polarization and $F_x(\mathbf{r}_2) = (3/2)|\alpha_1 E_{0y}|^2/a^4$ in s -polarization. The points (with the symbols + and $\times$) obtained with this approximation are shown in Fig. 2a and fit correctly the curves obtained without any approximation, as seen in this figure. Thus, they validate the approximation $\mathbf{T}(\mathbf{r}_i^k, \mathbf{r}_j^l, \omega)\alpha_j^l \ll 1$ previously done. Only when the spheres are very close to each other this approximation slightly departs from the exact calculation due to the increase of the free space susceptibility. In fact, this approximation assumes the dipole associated to the spheres as only due to the incident field, which is a good assumption when the polarizabilities are small, like for glass spheres. It is now easy to physically understand from Eq. (5) the reason of this either attractive or repulsive force. As the spheres are small, the scattering force is negligible²² and thus only the gradient force remains due to the interaction between the dipole associated to sphere B, and to the variation of the field created by sphere A at the position of sphere B. In p -polarization, the field due to sphere A at the position of sphere B and the dipole of sphere B are in phase, hence sphere B is pushed to the higher intensity region, namely towards sphere A. In s -polarization, as the field due to sphere A at the position of sphere B and the dipole of sphere B are in opposition of phase, sphere B is pushed to the lower intensity region, namely, far from sphere A. One can observe a similar effect in an atom mirror²³, or on a small silver particle in an evanescent field.¹⁶

On the other hand, in the far field we obtain, from the appendix, the force upon sphere B as: $F_x(\mathbf{r}_2) = |\alpha_1 E_{0x}|^2 k_0^2 \cos(k_0 a)/a^2$ in p -polarization and $F_x(\mathbf{r}_2) = -|\alpha_1 E_{0y}|^2 k_0^3 \sin(k_0 a)/(2a)$ in s -polarization, with $k_0 = 2\pi\sqrt{\epsilon_w}/\lambda$. The same explanation as before can be used for the sign of the force: following the phase relationship between the dipole and the field due to sphere A, the force is either positive or negative, hence, the oscillations of the force F_x take place with period $\lambda/\sqrt{\epsilon_w}$. The phase difference $\lambda/(4\sqrt{\epsilon_w})$ which appears in the far field between the oscillations of s and p polarization, comes from the difference between the derivative in the components xx and yy of the free space susceptibility. We observe that the force in p -polarization decreases faster than in s -polarization, this is due to the absence of a propagating field along the X -axis in the far field. The

magnitude of the force differs by a factor 10^4 between far field and near field. This is due to the strong interaction between the spheres through the evanescent waves.

We can make an analogy in the near field with molecular physics. If we look at the dipole moment of the two spheres, we compare our system of forces with the interaction between two molecules. In p -polarization, as the dipole moments are aligned and antisymmetric, they produce an attractive force analogous to that between two orbitals p_z , giving rise to a bonding state σ_u . In s -polarization, the dipole moments are parallel and symmetric, so we have antibonding states π_g^* , where $*$ means that the two spheres cannot be bound.




FIG. 3. Normalized force in the Z -direction on sphere B versus distance x between the centers of the spheres. Both spheres are of glass with $r = 10\text{nm}$, $\theta = 0^\circ$ and $\lambda = 632.8\text{nm}$ in vacuum. The full line corresponds to p -polarization, whereas the dashed line represents s -polarization.

We represent in Fig. 3 the force along the Z -direction. In this case the scattering force is predominant. The interaction between the spheres is now directly responsible for the oscillation of the force. Notice that when the spheres are far from each other, as the interaction between the spheres becomes weak when the distance increases, the force tends toward the scattering force upon one sphere due to the incident field. As this force is not responsible of optical binding, we are not going to discuss it further.




FIG. 4. Non-retarded approximation normalized force in the X -direction on sphere B versus distance x between the centers of the spheres. The sphere A is of silver and the sphere B is of glass with $r = 10\text{nm}$. $\lambda = 365\text{nm}$ (full line), 388.5nm (dotted line), and 600nm (dashed line). Without symbol: p -polarization; with symbol $+$: s -polarization.

More interesting is the case of two different small spheres, one (B) being dielectric and the other (A) being metallic (silver). The first fact easily observed from Eq. (4) pertains to the non retarded case, as the derivative of the free space susceptibility is real, the forces on spheres A and B are equal but of opposite sign to each other (this is no longer the case in the far field). In Fig. 4 we represent the force on sphere B at short distances from sphere A (in comparison with the wavelength), for $\lambda = 365\text{nm}$, 388.5nm , and 600nm . For p -polarization, we observe that the force at the wavelength $\lambda = 365\text{nm}$ has a rather strange behavior as it is positive, and only when the spheres are almost in contact, this force changes and becomes similar to those at the other wavelengths. With the approximation $\mathbf{T}(\mathbf{r}_i^k, \mathbf{r}_j^l, \omega)\alpha_j^l \ll 1$, the force in the non retarded approximation can be written

$$F_x(\mathbf{r}_2) = 1/2 \frac{\partial T_{ii}(\mathbf{r}_2, \mathbf{r}_1, \omega)}{\partial x} |E_{0i}|^2 \alpha_2 \Re(\alpha_1). \quad (6)$$

We observe that the sign of the force depends of $\Re(\alpha_1)$. When $\Re(\alpha_1) > 0$ ($\lambda = 600\text{nm}$), which is the common case, the dipole associated to the silver sphere is in phase with the applied field, so everything happens as for the dielectric sphere. Conversely, when $\Re(\alpha_1) < 0$ ($\lambda = 365\text{nm}$) the dipole is in opposition of phase with the applied field, and hence the force becomes positive. But when the spheres are almost in contact, the approximation $\mathbf{T}(\mathbf{r}_i^k, \mathbf{r}_j^l, \omega)\alpha_j^l \ll 1$ is no longer valid as shown when $\Re(\alpha_1) = 0$ ($\lambda = 388.5\text{nm}$), case in which the force is not null but negative. This is due to the fact that the polarizability of the silver sphere is large and hence the approximation is no longer valid for short distances. Physically, this represents the contribution of the metallic sphere to the electric field acting on the dielectric sphere,

which is larger than that of the incident field, hence the dipoles associated to the two spheres are in phase and the force is attractive. Notice that the change of sign occurs both at the plasmon resonance ($\Re(\varepsilon_1) \approx -2\varepsilon_w$) and when $\Re(\varepsilon_1) \approx \varepsilon_w$. When ε_1 is between these two values, the real part of the polarizability is negative. For a more complete discussion on this, one can see Ref. [16]. A similar reasoning can be done for s -polarization. If we now make the analogy previously done with the molecular orbitals, then depending on the wavelength in s -polarization, we shall obtain either antibonding states π_g^* , or bonding states π_u .

In the far field, for p -polarization, on using Eq. (4) and the approximation $\mathbf{T}(\mathbf{r}_i^k, \mathbf{r}_j^l, \omega)\alpha_j^l \ll 1$, we can write the force on the sphere B as

$$F_x(\mathbf{r}_2) = (\Re(\alpha_1^* \alpha_2) \cos(k_0 a) - \Im m(\alpha_1^* \alpha_2) \sin(k_0 a)) k_0^2 |E_{0x}|^2 / a^2 \quad (7)$$

and the force on sphere A as:

$$F_x(\mathbf{r}_1) = (-\Re(\alpha_1^* \alpha_2) \cos(k_0 a) - \Im m(\alpha_1^* \alpha_2) \sin(k_0 a)) k_0^2 |E_{0x}|^2 / a^2 \quad (8)$$

As the spheres are small, we can take only the gradient force as this is now the predominant one, then α_2 is real.¹⁵ Therefore, the forces on spheres A and B for the wavelength $\lambda = 600\text{nm}$, where $\Im m(\alpha_1)$ is weak, are opposite to each other like for two identical spheres. But at $\lambda = 388.5\text{nm}$, where $\Re(\alpha_1) = 0$, the forces on the two spheres are completely identical.




FIG. 5. Two glass spheres of radius $r=100\text{nm}$, $\theta = 0^\circ$, and $\lambda = 632.8\text{nm}$ in vacuum. The laser intensity of the incident light is $0.2\text{W}/\mu\text{m}^2$. Full line curves are for p -polarization. Dashed line curves correspond to s -polarization. a) Potential of sphere B normalized to $k_b T$ versus distance between the centers of the spheres. The height of the bars correspond to a normalized potential equal to 3. b) Force in the X -direction versus the distance between the spheres.

It should be remarked that if the laser intensity of the incident light is assumed to be $0.2\text{W}/\mu\text{m}^2$,²⁴ the optical forces for these small spheres are not strong enough to create an optical binding, since then the Brownian motion stays the dominant force. In this respect, the interest of the case of these small spheres is mainly the interpretative value it yields of the underlying physics. However, for larger radius in comparison to the wavelength, the forces become larger and so is the trapping potential. In Figs. 5b and 6b we plot the force along the X -axis for two dielectric spheres (glass) with radius $r = 100\text{nm}$, and 200nm , respectively. We observe that with the intensity used previously ($0.2\text{W}/\mu\text{m}^2$) the magnitude of the force is now enough to optically binding both spheres. We compute the potential energy of the optical trap by integration of the force (we take the potential energy null when the second sphere is at infinity). As the two spheres are identical, the potential energy is the same for both. The efficiency of the trapping force requires it to be larger than the force due to the Brownian motion, hence the depth of the potential wells of the trap should be larger than k_bT , T being the temperature of water, and k_b the Boltzmann constant. Considering $T = 290\text{K}$, then $k_bT = 4 \times 10^{-21}\text{J}$. We plot in Figs. 5a and 6a the potential normalized to the value k_bT . We adopt the criterion that the trap is efficient when the potential well is larger than $3k_bT$. Hence the bars plotted at the bottom of the wells in Figs. 5a and 6a correspond to the value 3. We see from Fig. 5 that for p -polarization the trap is not feasible except when the spheres are in contact. For s -polarization, we have three equilibrium positions spaced out by one wavelength. This behavior is explained by the previous results on small spheres, and in agreement with experiments.¹⁰ When the size of the sphere is close to one wavelength, we see from Fig. 6 that the depth of the potential well is larger than in the previous case. In p -polarization there is no possibility to stick the spheres together, but now we have one potential minimum of stable position. Thus, we observe that it is easier to trap particles when their radius are large, in agreement with the experiments of Burns et al.¹⁰.




FIG. 6. Same as Fig. 5 but with a radius $r = 200\text{nm}$.

Notice that the gravity force is $6.16 \times 10^{-5} \text{ pN}$ and $4.93 \times 10^{-4} \text{ pN}$ for the spheres of radii $r=100\text{nm}$ and 200nm , respectively. Another force that exists between the spheres is the Casimir-Polder force, F_c . To our knowledge, F_c has often been studied either between two plates or between a sphere and a plate,²⁵ but it has never been established between two spheres. In the non retarded approximation and the dipole approximation for the spheres, F_c is reduced to the dispersion force (London's force) which is inversely proportional to the seventh power of the distance between the dipoles (see for example Ref. [26]). Hence, only when the spheres are in contact, or at distances smaller than the wavelength from each other, this force might be of the same magnitude of the optical forces. However, the fast decay of this force at distances larger than the wavelength prevents it to perturb the optical trap.

In table 1 we give some examples on the limiting radius to get optical trapping for two identical spheres embedded in water (notice that for p polarization we do not mean optical trapping when the spheres are stucked in contact), i.e., the minimum radius to obtain one potential minimum of stable position for the two spheres using the same criterion as before (namely, $U > 3k_bT$).

spheres	glass ($\lambda = 632.8\text{nm}$)	silver ($\lambda = 394\text{nm}$)	silver ($\lambda = 314\text{nm}$)
limiting radius (p -pol.)	123 nm	33 nm	180 nm
limiting radius (s -pol.)	85 nm	21 nm	50 nm

TABLE I. Minimum radius to get one minimum position of the potential for two identical spheres. The following cases are addressed: glass sphere, silver sphere (both off and in plasmon resonance) The criterion of stability used is: the potential well depth must be larger than $3k_bT$.

We should remark that this is the limiting radius only to obtain the first stable position, if we want to get more stable positions, as in Ref.[10], the radius must be larger.

As mentioned before, the table shows that the optical trapping is easier for s polarization. For the silver sphere, the value $\lambda = 394\text{nm}$ corresponds to the plasmon reso-

nance, and that of $\lambda = 314\text{nm}$ is for a wavelength out of resonance. At the plasmon resonance the polarizability is largest, so it is easier to perform optical binding at this wavelength.




FIG. 7. The sphere A is of glass and the sphere B of silver with $r = 100\text{nm}$, $\lambda = 388.5\text{nm}$. The laser intensity of the incident light is $0.2\text{W}/\mu\text{m}^2$. Plot of the potential normalized to $k_b T$ for the two polarizations and the two spheres, since now this magnitude depends on the sphere material. We then observe that it is not possible to obtain a stable equilibrium since the potentials of the two spheres are now different. This result is explainable from the previous calculation on small spheres (of silver and glass), since then the forces were always opposed to each other at this wavelength. In fact, as the spheres are now large, the forces at this wavelength are not exactly in opposition, due to the larger scattering and absorbing force. Hence it is possible to obtain points where the potential of the two spheres is minimum. This happens when the forces on each sphere are the same and positive. In that case, the two spheres move in the direction of the positive X -axis while keeping constant the distance between them.

As a last instance, we now consider two spheres in free space, with radii $r = 100\text{nm}$, one being of glass and the other of silver, illuminated by a plane wave at $\lambda = 388.5\text{nm}$. We plot in Fig. 7 the potential energy for the two polarizations and the two spheres, since now this magnitude depends on the sphere material. We then observe that it is not possible to obtain a stable equilibrium since the potentials of the two spheres are now different. This result is explainable from the previous calculation on small spheres (of silver and glass), since then the forces were always opposed to each other at this wavelength. In fact, as the spheres are now large, the forces at this wavelength are not exactly in opposition, due to the larger scattering and absorbing force. Hence it is possible to obtain points where the potential of the two spheres is minimum. This happens when the forces on each sphere are the same and positive. In that case, the two spheres move in the direction of the positive X -axis while keeping constant the distance between them.

B. Particles in water on a dielectric flat surface

In this section we consider a flat dielectric surface upon which the spheres are suspended in water, as shown by Fig. 1. We compute the force along the X -axis on sphere B when both spheres are dielectric (glass), with $\theta = 0^\circ$ (Fig. 8). We now observe that both for large and small spheres the force has a behavior similar to that acting on

dielectric spheres isolated in water. When the spheres are in contact, the force on sphere B is the same as in absence of interface, whereas when the spheres are far from each other, this force is slightly smaller than without interface. This means that the optical binding is more difficult to perform when the spheres are on a surface than when they are far from interfaces. Also, there is a change in the period of oscillation due to interaction between the spheres via the light reflected by the surface. However, in the case when one of the spheres is metallic (silver), we observe the same behavior of the forces when the surface is present as without it. Then, as previously observed for dielectric spheres, there only appears a shift in the oscillation and magnitude of the forces.




FIG. 8. Force in the X -direction upon sphere B when the spheres are placed on a flat dielectric surface. $\theta = 0^\circ$, $\lambda = 632.8\text{nm}$. The full line represents the force in the presence of the surface, and the dashed line corresponds to the force computed without the interface. The curves with symbols + denote s -polarization, and those without symbol correspond to p -polarization. a) spheres of glass with $r = 10\text{nm}$. b) spheres of glass with $r = 100\text{nm}$.

In Fig. 9 we investigate the potential and optical force on two glass spheres in front of a dielectric surface, illuminated by total internal reflection ($\theta = 50^\circ$). Figure 9b shows that the force component along the X -axis always pushes the spheres in the direction of the wave vector component parallel to the surface. Hence, it is not possible to obtain a stable equilibrium with the two spheres remaining fixed. But if we compute the potential of the two spheres together (Fig. 9a), we observe some minima indicating that the system can acquire internal equilibrium, namely, the relative positions of the spheres can be kept fixed. Hence, when both spheres move impelled by the evanescent wave propagating along the surface, their velocity remains parallel to this surface, while the distance between them keeps some particular values given by the position of the potential minima (cf. Fig. 9a). Notice that the force on the second sphere (in both polarizations) has no oscillation, a very similar behavior

was observed by Okamoto et .*al.*²⁴. The computational prediction of similar collective movements in systems of more than two spheres will involve long computing times of their relative positions by potential energy minimization.

IV. CONCLUSION

We have studied the optical binding between two spheres embedded in water, either in presence or absence of a flat dielectric interface. We have presented results for different sizes and illumination conditions. Some of them agree with previous experiments¹⁰ for two identical spheres, however, when they are composed of different material, the force between them may have quite different behavior, depending on the light wavelength employed. In future work, it would be interesting to investigate the effect of light on several spheres in water in order to build up particle arrays. However, in this respect, the vertical force that pushes the spheres away from the substrate, constitutes a hindrance to this aim. This work shows, however, that this problem can be avoided by illuminating the system under total internal reflection at the substrate interface. Then the spheres will be stucked to the surface by the gradient force due to the transmitted evanescent wave. The horizontal force of this surface wave on the spheres that pushes them along the interface, can be compensated by means of a second counter-propagating evanescent wave, created by an additional beam. Notice, in addition, that if both surface waves are mutually coherent, the resulting standing wave pattern can introduce further structure in the resulting potential wells acting on the spheres.




FIG. 9. Fig. 9 Two glass spheres of radius $r=100\text{nm}$ in vacuum, in front of a flat dielectric surface. $\theta = 50^\circ$, $\lambda = 632.8\text{nm}$. The laser intensity of the incident light is $0.2\text{W}/\mu\text{m}^2$. Full line curves are for p -polarization. Dashed line curves correspond to s -polarization. a) Potential of interaction between the two spheres, normalized to $k_b T$, versus distance between the centers of the spheres. b) Force in the X -direction against distance between the spheres. The force curve on sphere A is without symbol, and the force on sphere B is with symbol +.

V. ACKNOWLEDGEMENTS

Work supported by the European Union and the Direction General de Investigacion Cientifica y Tecnica, grant PB98 0464.

APPENDIX A: DERIVATIVE OF THE FREE SPACE SUSCEPTIBILITY

The derivative of the free space susceptibility used in this paper is:

$$\frac{\partial}{\partial x} T_{xx}(x, x_0) = -\frac{6\mathbf{a}}{a^5}, \quad (\text{A1})$$

$$\frac{\partial}{\partial x} T_{yy}(x, x_0) = \frac{\partial}{\partial x} T_{zz}(x, x_0) = \frac{3\mathbf{a}}{a^5}, \quad (\text{A2})$$

in the non retarded case, and

$$\frac{\partial}{\partial x} T_{xx}(x, x_0, \omega) = \frac{2ak_0^2}{a^3} e^{ik_0 a}, \quad (\text{A3})$$

$$\frac{\partial}{\partial x} T_{yy}(x, x_0, \omega) = \frac{\partial}{\partial x} T_{zz}(x, x_0, \omega) = \frac{ia_0^3}{a^2} e^{ik_0 a}, \quad (\text{A4})$$

in the far field, where $\mathbf{a} = (x - x_0)$ and $a = |\mathbf{a}|$. x is the abscissa of the observation point, and x_0 that of dipole position.

-
- ¹ A. Ashkin, Phys. Rev. Lett. **24**, 156 (1970).
 - ² A. Ashkin, Phys. Rev. Lett. **25**, 1321 (1970).
 - ³ A. Ashkin, J. M. Dziedzic, J. E. Bjorkholm, and S. Chu, Opt. Lett. **11**, 288 (1986).
 - ⁴ A. R. Clapp, A. G. Ruta, and R. B. Dickinson, Rev. Sci. Instr. **70**, 2627 (1999).
 - ⁵ A. C. Dogarin and R. Rajagopalan, Langmuir **16**, 2770 (2000).
 - ⁶ A. Ashkin and J. M. Dziedzic, Science **235**, 1517 (1987).
 - ⁷ S. D. Collins, R. J. Baskin and D. G. Howitt, Appl. Opt. **38**, 6068 (1999).
 - ⁸ R. C. Gauthier and A. Frangioudakis, Appl. Opt. **39**, 26 (2000).

- ⁹ K. Taguchi, K. Atsuta, T. Nakata, and M. Ikeda, Opt. Comm. **176**, 43 (2000).
- ¹⁰ M. Burns, J.-M. Fournier, and J. Golovchenko, Phys. Rev. Lett. **63**, 1233 (1989).
- ¹¹ M. I. Antonoyiannakis, and J. B. Pendry, Phys. Rev. B **60** 2363 (1999).
- ¹² M. I. Antonoyiannakis, and J. B. Pendry, Europhys. Lett. **40** 613 (1997).
- ¹³ M. Bayer, T. Gutbrod, A. Forchel, T. L. Reinecke, P. A. Knipp, A. A. Dremin, V. D. Kulakovskii, and J. P. Reithmaier, Phys. Rev. Lett. **81** 2582 (1997).
- ¹⁴ P. C. Chaumet, and M. Nieto-Vesperinas, Phys. Rev. B. **61**, 14119 (2000).
- ¹⁵ P. C. Chaumet, and M. Nieto-Vesperinas, Opt. Lett. **25**, 1065 (2000).
- ¹⁶ P. C. Chaumet, and M. Nieto-Vesperinas, Phys. Rev. B. **62**, 11185 (2000).
- ¹⁷ Purcell and Pennypacker, Astrophys. J. **186**, 705 (1973).
- ¹⁸ J. A. Stratton, *Electromagnetic theory*, McGraw-Hill, New-York, 1941.
- ¹⁹ J. D. Jackson, *Classical Electrodynamics*, 2nd ed. (John Wiley, New York, 1975), p395.
- ²⁰ G. S. Agarwal, Phys. Rev. A **11**, 230 (1975); **12**, 1475 (1975).
- ²¹ A. Rahmani, P. C. Chaumet, F. de Fornel, and C. Girard, Phys. Rev. A **56**, 3245 (1997).
- ²² For a small sphere, the gradient and scattering forces are proportional to a^3 and a^6 , respectively.
- ²³ L. Cognet, V. Savalli, G.Zs.K. Horvath, D. Holleville, R. Marani, C. I. Westbrook, N. Westbrook, and A. Aspect, Phys. Rev. Lett. **81**, 5044 (1998), A. Landragin, J.-Y. Courtois, G. Labeyrie, N. Vansteenkiste, C. I. Westbrook, and A. Aspect, Phys. Rev. Lett. **77**, 1464 (1996).
- ²⁴ K. Okamoto and S. Kawata, Phys. Rev. Lett. **83**, 4534 (1999).
- ²⁵ G. L. Klimchitskaya, U. Mohideen, and V. M. Mostepanko, accepted at Phys. Rev. A.
- ²⁶ P. W. Milonni, Phys. Rev. A **53**, 3484 (1996).